



Acthar[®]GEL
(repository corticotropin injection) 80 U/mL

Acthar Referral Form

FAX: 1-877-937-2284

EMAIL: intake@supportandaccess.com

Please complete and email or fax toll-free

For questions, please call: 1-888-435-2284

Monday through Friday (8:00 AM to 9:00 PM ET)

Saturday (9:00 AM to 2:00 PM ET)

PRESCRIBER INSTRUCTIONS:

1. **Have your patient read page 3 (section 9): PATIENT AUTHORIZATION(S). Request that the patient sign the top section** to allow Acthar Patient Support to provide a complete level of support during the approval process. **If the patient would like to receive support, please have them sign the second section or provide consent at ActharConsent.com** to enroll in support and educational programs to receive additional information about their condition and treatment.
2. **Complete pages 1 and 2 of the Acthar Referral Form.**
3. **Email or fax the completed Acthar Referral Form along with clinical notes, any medically relevant documentation, and copies of both the front and back of your patient's medical and prescription benefit card(s) to 1-877-937-2284 or intake@supportandaccess.com.**
4. Acthar Patient Support will process the Acthar Referral Form and contact both you and your patient.
5. Prior authorization assistance will only be provided for indicated disease states. Medicare, Medicaid, and other federal or state healthcare program patients may be ineligible for certain other aspects of Acthar assistance programs.

PRESCRIBER SIGNATURE ON PAGE 1 AUTHORIZES PRESCRIPTION, CONSENT, AND STATEMENT OF MEDICAL NECESSITY

By signing page 1, I certify that Acthar[®] Gel is medically necessary for this patient and that I have reviewed this therapy with the patient and will be monitoring the patient's treatment. I verify that the patient and Prescriber information on this enrollment form was completed by me or at my direction and that the information contained herein is complete and accurate to the best of my knowledge.

I understand that I must comply with my practicing state's specific prescription requirements, such as e-prescribing, state-specific prescription form, fax language, etc. Noncompliance with state-specific requirements could result in outreach to me by the dispensing pharmacy.

I authorize United BioSource LLC ("UBC"), the current operator of Acthar Patient Support, and other designated operators of the Program, to act on my behalf for the limited purposes of transmitting this prescription to and received by the designated Specialty Pharmacy by any means under applicable law, including via a designated third party or other operator of the Program.

I understand that representatives from the Program or UBC may contact me or my patient for additional information relating to this prescription. I acknowledge and agree that this prescription may be sent to and received by the designated Specialty Pharmacy by any means under applicable law, including via a designated third party or other operator of the Program, and that no additional confirmation of receipt of prescription is required by the designated Specialty Pharmacy.

I request that company-funded Acthar Injection Training Services be arranged for my patient. I understand that Acthar Injection Training Services are available for multiple visits but are NOT a home health nursing service and that I or my patient may opt out of any nursing services by notifying the Acthar Patient Support Team by calling 1-800-435-2284. Patients can contact their Nurse Navigator at any time about injection training.

PATIENT INSTRUCTIONS:

Your Prescriber will submit the completed Acthar Referral Form to Acthar Patient Support. After we receive the form, we will call you so we can help you get your medicine. Please be on the lookout and answer calls from 1-800, 1-888, or blocked numbers. If you have any questions, please call **1-888-435-2284** Monday through Friday from 8 AM to 9 PM ET or Saturday from 9 AM to 2 PM ET.



Acthar[®]GEL

(repository corticotropin injection) 80 U/mL

FAX: 1-877-937-2284 or EMAIL: intake@supportandaccess.com

☐ SENT PRESCRIPTION DIRECTLY TO SPECIALTY PHARMACY.
PLEASE ENROLL PATIENT IN ACTHAR PATIENT SUPPORT.

PHARMACY NAME: _____

Acthar Referral Form

Please complete and email or fax toll-free
For questions, please call: 1-888-435-2284
Monday through Friday (8:00 AM to 9:00 PM ET)
Saturday (9:00 AM to 2:00 PM ET)

1. PATIENT INFORMATION

Patient has been notified of referral ☐ YES ☐ NO

PATIENT FIRST NAME	MIDDLE INITIAL	LAST NAME	DATE OF BIRTH	GENDER
HOME ADDRESS			CITY	STATE ZIP
SHIPPING ADDRESS (IF NOT HOME)	CARE OF (IF NOT ADDRESSED TO PATIENT)	CITY	STATE	ZIP
HOME PHONE	MOBILE PHONE	ALTERNATE PHONE	BEST TIME TO CALL	
EMAIL ADDRESS	PREFERRED LANGUAGE IF NOT ENGLISH			
ALTERNATIVE CONTACT NAME	TELEPHONE	EMAIL	RELATIONSHIP TO PATIENT	

2. INSURANCE INFORMATION (Please include copies of front and back of all medical and prescription insurance cards)

PHARMACY BENEFITS	SUBSCRIBER ID #	GROUP #	TEL #
PRIMARY MEDICAL INSURANCE	SUBSCRIBER ID #	GROUP #	TEL #

3. PRESCRIBER INFORMATION SPECIALTY: ☐ NEUROLOGY ☐ OTHER (Please indicate on line 2 below)

PRESCRIBER FIRST NAME	MIDDLE INITIAL	LAST NAME	NPI #	STATE LICENSE #
OFFICE / CLINIC / INSTITUTION NAME	TELEPHONE	FAX	OTHER SPECIALTY	
ADDRESS	CITY	STATE	ZIP	
OFFICE CONTACT NAME	CONTACT TELEPHONE	CONTACT MOBILE PHONE	CONTACT EMAIL ADDRESS	

4. PRESCRIPTION: ACTHAR[®] GEL NDC# 63004-8710-1 5 mL multidose vial containing 80 USP units per mL inj

4A. ICD-10 CODE: (REQUIRED):

☐ MULTIPLE SCLEROSIS G35 ☐ OTHER OPTIC NEURITIS H46.8

NOTE: PROVIDED TO THE LEFT ARE THE **2 MOST COMMON CODES**; FOR OTHER PRIMARY DIAGNOSIS CODES, PLEASE SEE PAGE 2, SECTION 6

4B. SELECT AN FDA RECOMMENDED DOSE OR OTHER DOSE

DOSE: ☐ 80 UNITS (1 mL) DAILY ☐ OTHER (Specify Units or mL): _____

FREQUENCY/DURATION: ☐ 10 DAYS (2 vials) ☐ 15 DAYS (3 vials) ☐ 20 DAYS (4 vials)

☐ OTHER: _____

ROUTE OF ADMINISTRATION WILL BE SUBCUTANEOUS UNLESS INTRAMUSCULAR IS SPECIFIED:

☐ INTRAMUSCULAR

QUANTITY OF 5 mL MULTIDOSE VIALS*: _____ REFILLS*: _____

4C. TAPER INSTRUCTIONS (Attach taper schedule and provide additional instructions below, if applicable)

4D. ALLERGIES ☐ NKDA - No known drug allergies (Additional space provided on pg 2)

4E. SUPPLIES PHARMACY TO SUPPLY FOLLOWING UNLESS "OTHER" IS SPECIFIED:

- SYRINGE: 1 ML (80 Unit dose) 3 ML (120 Unit dose)
- NEEDLE FOR DRAWING: 20 G
- NEEDLE FOR INJECTION:
25 G, 5/8" (SUBCUTANEOUS) OR
25 G, 1" (INTRAMUSCULAR)—if box checked in 4B
- SHARPS CONTAINER

PHARMACY TO DISPENSE SUFFICIENT SUPPLIES TO COMPLETE COURSE OF THERAPY. PHARMACIST MAY ELECT TO DISPENSE ALTERNATE SUPPLIES AS NECESSARY.

☐ OTHER:

5. HISTORY OF CORTICOSTEROID USE (PLEASE ADD DETAILS IN SECTION 6 OR 7 ON PAGE 2).

PLEASE CHECK ALL THAT APPLY:

A corticosteroid **was** tried with the following response(s) - please provide specific evidence for reaction to or intolerance of corticosteroids below.

- ☐ Corticosteroid use failed, but same response not expected with Acthar
- ☐ Patient hypersensitive or allergic to corticosteroids
- ☐ Patient intolerant of corticosteroids

as evidenced by:

A corticosteroid **was not** tried due to the following reason(s) - please provide specific evidence of contraindication or intolerance below.

- ☐ Patient is homebound
- ☐ Corticosteroid use is contraindicated for this patient
- ☐ Intravenous access is not possible for this patient
- ☐ Patient has known intolerance to corticosteroids

as evidenced by:

OPT OUT ONLY - ACTHAR INJECTION TRAINING SERVICES

☐ By checking here, I request to opt out of Acthar Injection Training Services for my patient.

PRESCRIBER SIGNATURE: Please sign only ONE LINE below (by signing below you are agreeing to the Prescriber Consent section on the cover page of this document)

Brand Medically Necessary / Do Not Substitute / No Substitution / DAW / May Not Substitute

May Substitute / Product Selection Permitted / Substitution Permissible

☒

DISPENSE AS WRITTEN

DATE

OR ☒

SUBSTITUTIONS ALLOWED

DATE

Prescriber signature required for consent and to validate prescriptions. Prescriber attests that this is her/his signature. NO STAMPS. By signing, Prescriber certifies that the above is medically necessary.

ATTN: New York and Iowa providers, please submit electronic prescription. CA, MA, NC & PR: Interchange is mandated unless Prescriber writes the words "No Substitution"

MS

Acthar[®]GEL
(repository corticotropin injection) 80 U/mL

Patient Name: _____ Date of Birth: _____

6. DIAGNOSIS AND MEDICAL INFORMATION

DIAGNOSIS CODES: BELOW IS A LIST OF THE MOST COMMON CODES. THESE CODES HAVE BEEN PROVIDED FOR CONVENIENCE ONLY. THESE ARE NOT ALL POSSIBLE DIAGNOSIS CODES, AND NOT INTENDED TO INFLUENCE A DIAGNOSIS.

Please provide as much information as possible that corresponds with the patient's diagnosis. You may also write in the patient's diagnosis in the "OTHER DIAGNOSIS" section.

☐ **Multiple sclerosis** (Must check one)

Is Acthar to be used to treat an acute exacerbation?

☐ Yes☐ No, other: _____

Onset of acute exacerbation Date: _____

☐ **MULTIPLE SCLEROSIS**
G35☐ **OTHER OPTIC NEURITIS**
H46.8☐ **OTHER DERMATOPOLYMYOSITIS WITH MYOPATHY**
M33.12☐ **OTHER DERMATOPOLYMYOSITIS WITHOUT MYOPATHY**
M33.13☐ **POLYMYOSITIS, ORGAN INVOLVEMENT UNSPECIFIED**
M33.20☐ **POLYMYOSITIS WITH MYOPATHY**
M33.22☐ **DERMATOPOLYMYOSITIS, UNSPECIFIED WITHOUT MYOPATHY**
M33.93☐ **OTHER DIAGNOSIS:****IMPORTANT!****7. RELEVANT TREATMENT HISTORY (INCLUDING RECENT CORTICOSTEROID HISTORY. ATTACH ADDITIONAL CASE NOTES AS NECESSARY.)**

Therapy Name	Dose	Start Date	Stop Date (if applicable)	Explain Outcome With Detail (eg, type of outcome)

8. OTHER RELEVANT CLINICAL INFORMATION (INCLUDING ALLERGIES)☐ NKDA - No known drug allergies**PRESCRIBER SIGNATURE: REQUIRED FOR DOCUMENTATION**

I verify that the patient and Prescriber information on this enrollment form was completed by me or at my direction and that the information contained herein is complete and accurate to the best of my knowledge. I certify that my patient has agreed in writing to be contacted by Program administrators or UBC and be furnished with Program or other information or materials.



NAME

X

SIGNATURE

DATE

MS

Acthar[®]GEL
(repository corticotropin injection) 80 U/mL

Patient Name: _____ Date of Birth: _____

9. PATIENT AUTHORIZATION(S)

Patient Consent to allow Acthar Patient Support Team to work together with your insurance provider, pharmacy, advocacy organization and others to provide support on your behalf.

By signing this authorization, I authorize my physician(s), my health insurance company and my pharmacy providers (collectively, "Designated Parties") to use, disclose, and redisclose to Mallinckrodt ARD LLC ("Mallinckrodt"), the distributor of Acthar, and its agents, authorized designees and contractors, including Mallinckrodt reimbursement support personnel and United BioSource LLC ("UBC") or any other operator of Acthar Patient Support on behalf of Mallinckrodt (collectively, "Manufacturer Parties"), health information relating to my medical condition, treatment and insurance coverage (my "Health Information") in order for them to (1) provide certain services to me, including reimbursement and coverage support, patient assistance and access programs, medication shipment tracking, and home injection training, (2) provide me with support services and information associated with my Acthar therapy, (3) serve internal business purposes, such as marketing research, internal financial reporting and operational purposes, and (4) carry out the Manufacturer Parties' respective legal responsibilities.

Once my Health Information has been disclosed to Manufacturer Parties, I understand that it may be redisclosed by them and no longer protected by federal and state privacy laws. However, Manufacturer Parties agree to protect my Health Information by using and disclosing it only for the purposes detailed in this authorization or as permitted or required by law.

I understand that I may refuse to sign this authorization and that my physician and pharmacy will not condition my treatment on my agreement to sign this authorization form, and my health plan or health insurance company will not condition payment for my treatment, insurance enrollment or eligibility for insurance benefits on my agreement to sign this authorization form. I understand that my pharmacies and other Designated Parties may receive payment in connection with the disclosure of my Health Information as provided in this authorization. I understand that I am entitled to receive a copy of this authorization after I sign it.

I may revoke (withdraw) this authorization at any time by mailing a letter to Acthar Patient Support, 680 Century Point, Lake Mary, FL 32746. Revoking this authorization will end further disclosure of my Health Information to Manufacturer Parties by my pharmacy, physicians, and health insurance company when they receive a copy of the revocation, but it will not apply to information they have already disclosed to Manufacturer Parties based on this authorization. I also know I may cancel my enrollment in a patient support program at any time in writing by contacting Mallinckrodt via fax at 1-877-937-2284 or by calling Acthar Patient Support at 1-888-435-2284. This authorization is in effect for 5 years unless a shorter period is provided for by state law (MARYLAND HEALTHCARE PROVIDERS, under Maryland Code HG § 4-303(b)(4) this authorization expires ONE YEAR from the date of signature) or until the conclusion of any ongoing coverage support, whichever is longer, once I have signed it unless I cancel it before then.

THIS SECTION MUST BE COMPLETED IN ITS ENTIRETY, INCLUDING DATE



X

PATIENT NAME OR LEGAL REPRESENTATIVE _____ PATIENT OR LEGAL REPRESENTATIVE SIGNATURE _____ IF LEGAL REPRESENTATIVE, RELATIONSHIP TO PATIENT _____ DATE _____

Patient Consent to receive additional information from Mallinckrodt such as education on your disease and Acthar.

I authorize Mallinckrodt and its partners to use, disclose, and/or transfer the personal information I supply (1) to contact me and provide me with informational and marketing materials and clinical trial opportunities related to my condition or treatment by any means of communication, including but not limited to text, email, mail, or telephone; (2) to help Mallinckrodt improve, develop, and evaluate products, services, materials, and programs related to my condition or treatment; (3) to enroll me in and provide me with Acthar-related programs and services that I may select or refuse at any time; (4) to disclose my enrollment and use of these services to my prescriber and insurers; and (5) to use my information that cannot identify me for scientific and market research. This authorization will remain in effect until I cancel it, which I may do at any time in writing by contacting Mallinckrodt via fax at 1-877-937-2284 or by calling Acthar Patient Support at 1-888-435-2284. I may request a copy of this signed authorization.

THIS SECTION MUST BE COMPLETED IN ITS ENTIRETY, INCLUDING DATE



X

PATIENT NAME OR LEGAL REPRESENTATIVE _____ PATIENT OR LEGAL REPRESENTATIVE SIGNATURE _____ IF LEGAL REPRESENTATIVE, RELATIONSHIP TO PATIENT _____ DATE _____

Scan the QR Code below to save the Acthar Patient Support phone number to your mobile device's contacts (see steps below).



STEP 1



Open the camera on your mobile device

STEP 2



Hold your camera over the QR code to scan

STEP 3



Save your Acthar Patient Support Team information to your contacts

If patient is not present to sign the form, send them to

ActharConsent.com

and have them sign electronically.



Acthar[®] GEL
(repository corticotropin injection) 80 U/mL

Acthar Patient Support

TEL: 1-888-435-2284

FAX: 1-877-937-2284

IMPORTANT SAFETY INFORMATION

Contraindications

Acthar is contraindicated:

- For intravenous administration
- In infants under 2 years of age who have suspected congenital infections
- With concomitant administration of live or live attenuated vaccines in patients receiving immunosuppressive doses of Acthar
- In patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, history of or the presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, primary adrenocortical insufficiency, adrenocortical hyperfunction, or sensitivity to proteins of porcine origin

Warnings and Precautions

- The adverse effects of Acthar are related primarily to its steroidogenic effects
- Acthar may increase susceptibility to new infection or reactivation of latent infections
- Suppression of the hypothalamic-pituitary-adrenal (HPA) axis may occur following prolonged therapy with the potential for adrenal insufficiency after withdrawal of the medication. Adrenal insufficiency may be minimized by tapering of the dose when discontinuing treatment. During recovery of the adrenal gland patients should be protected from the stress (e.g., trauma or surgery) by the use of corticosteroids. Monitor patients for effects of HPA axis suppression after stopping treatment
- Cushing's syndrome may occur during therapy but generally resolves after therapy is stopped. Monitor patients for signs and symptoms
- Acthar can cause elevation of blood pressure, salt and water retention, and hypokalemia. Monitor blood pressure and sodium and potassium levels
- Acthar often acts by masking symptoms of other diseases/disorders. Monitor patients carefully during and for a period following discontinuation of therapy
- Acthar can cause gastrointestinal (GI) bleeding and gastric ulcer. There is also an increased risk for perforation in patients with certain GI disorders. Monitor for signs of perforation and bleeding
- Acthar may be associated with central nervous system effects ranging from euphoria, insomnia, irritability, mood swings, personality changes, and severe depression to psychosis. Existing conditions may be aggravated
- Patients with comorbid disease may have that disease worsened. Caution should be used when prescribing Acthar in patients with diabetes and myasthenia gravis
- Prolonged use of Acthar may produce cataracts, glaucoma, and secondary ocular infections. Monitor for signs and symptoms
- Acthar is immunogenic and prolonged administration of Acthar may increase the risk of hypersensitivity reactions. Cases of anaphylaxis have been reported in the postmarketing setting. Neutralizing antibodies with chronic administration may lead to loss of endogenous ACTH and Acthar activity
- There may be an enhanced effect in patients with hypothyroidism and in those with cirrhosis of the liver
- Long-term use may have negative effects on growth and physical development in children. Monitor pediatric patients
- Decrease in bone density may occur. Bone density should be monitored in patients on long-term therapy

Adverse Reactions

- Commonly reported postmarketing adverse reactions for Acthar include injection site reaction, asthenic conditions (including fatigue, malaise, asthenia, and lethargy), fluid retention (including peripheral swelling), insomnia, headache, and blood glucose increased
- The most common adverse reactions for the treatment of infantile spasms (IS) are increased risk of infections, convulsions, hypertension, irritability, and pyrexia. Some patients with IS progress to other forms of seizures; IS sometimes masks these seizures, which may become visible once the clinical spasms from IS resolve

Pregnancy

- Acthar may cause fetal harm when administered to a pregnant woman

Please see accompanying full Prescribing Information for additional Important Safety Information or visit <https://www.actharhcp.com/Static/pdf/Acthar-PI.pdf>.

INDICATION AND USAGE

Acthar Gel is indicated for the treatment of acute exacerbations of multiple sclerosis in adults. Controlled clinical trials have shown Acthar to be effective in speeding the resolution of acute exacerbations of multiple sclerosis. However, there is no evidence that it affects the ultimate outcome or natural history of the disease.